Among the painkillers(4)	acute ostry
over the counter, paracetamol is very(5)	analgesic lek przeciwbólowy
thanks to its(6) safety and few	anti-inflammatory
interactions with other medicines. It does not, however, reduce	przeciwzapalny
inflammation which is often the cause of pain. Non-steroidal	aspirin ['asp(ə)rɪn] aspiryna
(7) drugs (NSAIDs) like aspirin,	codeine [ˈkəʊdiːn] kodeina
ibuprofen, or diclofenac reduce both pain and inflammation, but not	contraindication
everybody can take them. Firstly, they have many contraindications	przeciwwskazanie
and secondly, they interact with some groups of medication. Their	dependence uzależnienie
(8) use in healthy adults is deemed	diclofenac [dar'kləʊfənak]
(9), however.	diverse zróżnicowany
Opioids like tramadol, codeine, or morphine are	general ogólny
(10) analgesics, but because of their many	inflammation zapalenie
(11) effects they are prescription-only	morphine [ˈmɔːfiːn] morfina
medication. Their(12) use can cause	relief ulga
physical dependence.	tramadol ['tramədol] tramadol

19 Names of medicines in different languages, even if written in a similar or identical way, may have different pronunciation. Fill in the table.

	Medicine	Number of syllables pronounced	Syllable stressed (e.g. 1st, 2nd, etc.)
1	paracetamol		
2	aspirin		
3	ibuprofen		
4	diclofenac		
5	tramadol		
6	codeine		
7	morphine		

20 Fill in the gaps. The first letters of the missing words have already been written.

Ι.	Suppositories are a rectal d form.		
2.	D the tablet in water and drink the solution.		
3.	C are easier to swallow than round tablets.		
4.	. Do you prefer tablets or ndrops?		
5.	Place the tablet under the t		
6.	S tablet should split easily.		
7.	Non-steroidal anti-i		
8.	Paracetamol tablets are available over the c		
9.	A dosage form consists of a substances and e like bulking agents, lubricants, solvents, or preservatives.		

10 Read the dialogue and answer the questions.

- 1. How does the pharmacist make sure that she is giving the medication to the right person?
- 2. How many levothyroxine tablets has Ms Bowden been prescribed?
- 3. Why was she prescribed hormone replacement therapy?
- 4. What is the dosage form of the HRT?
- 5. How should it be used?

'Prescription for Ms Bowden!'

'Here, that's me.'

'Can you confirm your full name and address, please?'

'It's Constance Bowden, 12 Willow Road.'

'That's right. Ms Bowden, we don't have the full amount of your levothyroxine tablets. Is it OK if we owe you some?'

'How many have you got?'

'Fifty-six. We would still owe you a pack of twenty-eight.'

'That's fine. When can I collect it?'

'I've already ordered it so the tablets should be here tomorrow. You can collect them anytime within six months, just bring this owing slip with you. Another thing is, our records say you've been prescribed HRT for the first time.'

'Yes, Dr Patel wanted me to try it for my night sweats.'

'I just want to make sure you know how to use it.'

'I was told to apply one twice a week.'

'That's right. Stick it somewhere on the trunk below the waist. Don't use the same spot twice in a row.'

'What happens when I take a shower?'

'Just leave it on.'

11) Fill in each gap with two words so that the second sentence has the same meaning as the first one.

consecutive kolejny

levothyroxine

trunk tułów

[ˌlivəʊθʌɪˈrɒkˌsin/ ˌlivəʊθʌɪˈrɒksən] lewotyroksyna

records dokumentacja

l.	we have only 28	out of 56 tablets you need.	
	We are 28 tablets	the required	amount.

2. Is it OK for you to collect the rest later?

Do collecting the rest later?

3. Our records say you take aspirin tablets.

..... our records, you take aspirin tablets.

4. Leave it on.

Don't take

5. You've been prescribed it for the first time.

You've it before.

16 Explain to your patients how they should take their medication.

1. Simvastatin 40 mg

to lower cholesterol levels

Mitte: 28 tablets

2.

Levothyroxine 100 μg

1 om ac

for underactive thyroid

Mitte: 28 tablets

3.

Simethicone 125 mg

1 qds prn

to reduce bloating

Mitte: 28 caps

4.

Aspirin 75 mg 1 om cc

to thin the blood

Mitte: 28 dispersible tablets

5.

Ranitidine 150 mg tabs 56

1 bd

for acid reflux

6.

Glycerin 4 g

1 prn

to relieve constipation
Mitte: 12 suppositories

7.

Clotrimazole 200 mg

1 nocte

for vaginal thrush Mitte: 3 pessaries

8.

Ramipril 5 mg 28 caps

1 od

to lower blood pressure

clotrimazole [kləʊˈtrʌɪməˌzəʊl/kləʊˈtrʌɪməˌzəːl] klotrymazol

glycerin ['glɪs(ə)riːn/'glɪs(ə)rɪn] gliceryna

ramipril [rəˈmɪprəl/raˈmɪprəl] ramipryl

simethicone [sɪˈmɛθɪˌəʊn/ sʌɪˈmɛθɪˌəʊn] symetykon, simetikon

simvastatin ['simvə,stat(ə)n/
 ,simvə'stat(ə)n] symwastatyna

17 Calculate how many tablets/millilitres will be needed for 7 days' supply, provided the medication is taken according to prescriber's directions.

	Directions	Required quantity
1	1 om ac	
2	2 od pc	
3	1 bd	
4	2 tds	
5	1 qds	
6	15 ml bd	
7	10 ml nocte	

Read the words aloud. Underline those which have the sound AI (as in ice) and circle those that have the sound EI (as in make). Some words do not have either sound.

thyroid, label, strip, enzyme, vial, clinical, rectal, nasal, cystitis, insulin, diabetes, interaction, syringe, formulation, dispense, antibiotic, dissolve, bladder, subcutaneous, item, urea, prescribe, ion

automatic pipette pipeta

automatyczna beaker zlewka bulb pipette filler gruszka do napełniania pipety burette biureta dropper kroplomierz Erlenmeyer/ conical flask erlenmajerka, kolba stożkowa evaporating dish parownica funnel lejek glass rod bagietka

IN A LAB

Label the laboratory equipment.















2)



graduated cylinder cylinder miarowy measuring

pipette pipeta wielomiarowa

mortar and pestle moździerz i pistel

spot plate płytka porcelanowa z wgłębieniami

test tube probówka test tube rack

stojak na probówki volumetric flask

kolba miarowa volumetric

pipette pipeta jednomiarowa

wash bottle tryskawka

















16)

SPEAKING ACTIVITIES

UNIT 2, EX. 17

Student B (patient)

Answer your partner's questions according to information given below. DO NOT volunteer any information unless asked. If asked about something not mentioned below, make up the answer.

When your partner finishes interviewing you, check together if there were any pieces of information he or she forgot to ask about.

Location of pain: head (headache), especially forehead

Intensity: moderate *Type*: throbbing

Duration: since yesterday
Other symptoms: none
Probable cause: not known

Previous experience: you have such headaches two or three times a year

How treated before: aspirin helped with the pain

UNIT 3, EX. 15

Student A (pharmacist)

To reduce the fever, recommend paracetamol or ibuprofen. Let the customer choose between tablets, capsules, effervescent tablets, suspension, and suppositories (paracetamol), tablets, capsules, and suspension (ibuprofen). You can offer liquid dosage forms in different flavours.

To relieve the sore throat, offer lozenges or a throat spray with antiseptic and anaesthetic agents.